


Alcohol Indicators for Wellingborough – Local Alcohol Profiles for England, March 2016

Background

Public Health England produce a range of indicators to evidence the impact of alcohol across the country, which are published at <http://fingertips.phe.org.uk/profile/local-alcohol-profiles>.

The majority of these indicators measure hospital admissions and deaths caused by alcohol consumption. Alcohol-specific indicators consider illnesses that are wholly attributable to alcohol consumption, such as alcoholic liver disease and alcohol poisoning, while alcohol-related indicators also consider illnesses in which alcohol may be a contributory factor, such as hypertension and certain cancers.

Where alcohol is one of several risk factors, Alcohol Attributable Fractions are used to estimate the likelihood of the death/hospital admission being linked to alcohol based on diagnoses, patient age and gender. The sum total of all these probabilities can be used as proxy estimates of the numbers of deaths/hospital admissions were related to alcohol. For example:



There are two measures of alcohol-related admissions:

- A broad measure that considers all diagnosis codes for each admission
- A narrow measure that considers only the primary diagnosis and external cause codes (e.g. assault)

The broad measure provides an overall estimate of the wider impact of alcohol on health services, while the narrow measure was introduced by Public Health England as an indicator that is less sensitive to coding practices and more responsive to changes brought about by local initiatives.

Both measures can be further broken down to reflect the specific causes of admission, such as liver disease and cardiovascular disease.

All admission and mortality indicators are expressed as directly standardised rates (DSR) per 100,000 population in order to be able to compare different geographies and timeframes.

Wellingborough

Summary

- Wellingborough's broad and narrow rates of alcohol-related admission episodes are similar to those for Northamptonshire.
- Wellingborough has the highest rates of alcohol-specific mortality in its CIPFA comparator group. The rates for females and persons are significantly worse than those for England and the East Midlands.
- There has been a significant increase in Wellingborough's rate of mortality from chronic liver disease between 2013 and 2014.
- Wellingborough had the second highest rate of alcohol-related road traffic accidents in its CIPFA comparator group and is significantly worse than the England average.

Indicators where Wellingborough's rate is significantly better than the England average

RAG ratings - ■ = significantly better, ■ = no significant difference, ■ = significantly worse, □ = not measured

Indicator (RAG against East Midlands benchmark)	Latest Period	Wellingborough	East Midlands	England	Local DOT Since Last Period
Alcohol-specific hospital admission - Persons	2013/14	268.4	317.9	373.8	↑
Alcohol-specific hospital admission - Male	2013/14	373.5	427.8	515.1	↑
Alcohol-specific hospital admission - Female	2013/14	174.6	213.5	241.3	↑
Alcohol-related hospital admission (Broad) - Persons	2013/14	1,142.6	1,134.4	1,253.0	↑
Alcohol-related hospital admission (Broad) - Male	2013/14	1,537.4	1,541.6	1,714.6	↑
Admission episodes for alcohol-related conditions (Broad) - Persons	2013/14	1,903.9	1,934.1	2,111.2	↑
Admission episodes for alcohol-related conditions (Broad) - Male	2013/14	2,565.8	2,636.8	2,916.8	↑
Admission episodes for alcohol-related conditions (Narrow) - Male	2013/14	716.0	857.9	835.3	↓
Claimants of benefits due to alcoholism	2014	88.1	106.4	131.0	N/A
Admission episodes for alcohol-related malignant neoplasm conditions (Broad) - Male	2013/14	131.7	214.6	177.0	↓
Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Broad) - Persons	2013/14	258.4	326.6	393.9	↑

Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Broad) - Male	2013/14	362.0	471.6	579.2	↑
Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Broad) - Female	2013/14	165.3	187.4	218.2	↑
Admission episodes for alcohol-related alcoholic liver disease condition (Broad) - Female	2013/14	30.3	54.8	65.8	↑
Admission episodes for alcohol-related malignant neoplasm conditions (Narrow) - Male	2013/14	103.8	187.1	154.7	↓
Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Narrow) - Persons	2013/14	62.4	80.6	87.3	↑
Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Narrow) - Male	2013/14	84.1	113.3	124.5	↑
Admission episodes for alcohol-related intentional self-poisoning by and exposure to alcohol condition (Narrow) - Female	2013/14	37.7	59.1	65.4	↑

Indicators where Wellingborough's rate is **not significantly different** to the England average

RAG ratings - ■ = significantly better, ■ = no significant difference, ■ = significantly worse, □ = not measured

Indicator (RAG against East Midlands benchmark)	Latest Period	Wellingborough	East Midlands	England	Local DOT Since Last Period
Months of life lost due to alcohol - Male	2012 - 14	13.9	12.0	12.0	↓
Months of life lost due to alcohol - Female	2012 - 14	8.8	5.8	5.6	↑
Alcohol-specific mortality - Male	2012 - 14	22.1	15.9	11.6	↓
Mortality from chronic liver disease - Persons	2012 - 14	15.1	11.6	16.1	↑
Alcohol-related mortality - Persons	2014	49.4	45.4	7.4	↓
Alcohol-related mortality - Male	2014	68.1	64.7	11.5	↓
Alcohol-related mortality - Female	2014	34.2	28.9	45.5	↑
Alcohol-specific hospital admission - under 18s	2011/12 - 13/14	29.1	33.8	65.4	↓
Alcohol-related hospital admission (Broad) - Female	2013/14	820.3	784.3	28.8	↑
Alcohol-related hospital admission (Narrow) - Persons	2013/14	407.6	439.6	40.1	↑

Alcohol-related hospital admission (Narrow) - Male	2013/14	536.0	584.2	858.5	↑
Alcohol-related hospital admission (Narrow) - Female	2013/14	298.5	310.3	443.7	↑
Admission episodes for alcohol-related conditions (Broad) - Female	2013/14	1,359.7	1,332.5	593.7	↑
Admission episodes for alcohol-related conditions (Narrow) - Persons	2013/14	591.7	673.9	310.3	↓
Admission episodes for alcohol-related conditions (Narrow) - Female	2013/14	487.6	508.0	645.1	↑
Admission episodes for alcohol-related malignant neoplasm conditions (Broad) - Persons	2013/14	175.9	212.8	835.3	↓
Admission episodes for alcohol-related malignant neoplasm conditions (Broad) - Female	2013/14	221.1	213.0	474.8	↑
Admission episodes for alcohol-related cardiovascular disease conditions (Broad) - Persons	2013/14	1,019.6	929.8	175.8	↑
Admission episodes for alcohol-related cardiovascular disease conditions (Broad) - Male	2013/14	1,459.5	1,349.3	177.0	↑
Admission episodes for alcohol-related cardiovascular disease conditions (Broad) - Female	2013/14	674.6	594.2	176.5	↑
Admission episodes for alcohol-related alcoholic liver disease condition (Broad) - Persons	2013/14	105.8	83.8	1,049.3	↑
Admission episodes for alcohol-related alcoholic liver disease condition (Broad) - Male	2013/14	183.9	114.1	1,523.8	↑
Admission episodes for alcohol-related malignant neoplasm conditions (Narrow) - Persons	2013/14	148.8	184.1	672.9	↓
Admission episodes for alcohol-related unintentional injuries conditions (Narrow) - Persons	2013/14	138.2	152.5	105.3	↑
Admission episodes for alcohol-related unintentional injuries conditions (Narrow) - Male	2013/14	205.1	229.0	147.1	↑
Admission episodes for alcohol-related unintentional injuries conditions (Narrow) - Female	2013/14	77.2	81.8	150.7	↑
Admission episodes for alcohol-related mental and behavioural disorders due to use of alcohol condition (Narrow) - Female	2013/14	41.6	48.8	154.7	↑
Admission episodes for alcohol-related intentional self-poisoning by and exposure to alcohol condition (Narrow) - Persons	2013/14	42.4	51.7	149.3	↑
Admission episodes for alcohol-related intentional self-poisoning by and exposure to alcohol condition (Narrow) - Male	2013/14	48.8	44.3	141.8	↓

Indicators where Wellingborough's rate is significantly worse than the England average


RAG ratings - ■ = significantly better, ■ = no significant difference, ■ = significantly worse, = not measured

Indicator (RAG against East Midlands benchmark)	Latest Period	Wellingborough	East Midlands	England	Local DOT Since Last Period
Alcohol-specific mortality - Persons	2013/14	18.3	11.6	11.6	↑
Alcohol-specific mortality - Female	2013/14	14.7	7.4	7.4	↑
Admission episodes for alcohol-related malignant neoplasm conditions (Narrow) - Female	2013/14	194.2	183.5	149.3	↓
Alcohol Related Road Traffic Accidents	2012 - 14	49.2	33.7	26.4	N/A


Indicators where Wellingborough is an outlier its CIPFA comparator group – top 20%

--- = England average


Admission episodes for alcohol-related conditions (Narrow) - Female




Admission episodes for alcohol-related conditions (Narrow) - Male




Admission episodes for alcohol-related conditions (Narrow) - Persons




Admission episodes for alcohol-related malignant neoplasm conditions (Narrow) - Persons




Admission episodes for alcohol-related malignant neoplasm conditions (Narrow) - Male




Admission episodes for alcohol-related intentional self-poisoning by and exposure to alcohol condition (Narrow) - Female




Admission episodes for alcohol-related intentional self-poisoning by and exposure to alcohol condition (Narrow) - Persons




Admission episodes for alcohol-related malignant neoplasm conditions (Broad) - Male




Alcohol-specific hospital admission - Female



Admission episodes for alcohol-related malignant neoplasm conditions (Broad) - Persons




Admission episodes for alcohol-related alcoholic liver disease condition (Broad) - Female




Indicators where Wellingborough is an outlier its CIPFA comparator group – **bottom 20%**


Months of life lost due to alcohol - Female




Months of life lost due to alcohol - Male




Admission episodes for alcohol-related unintentional injuries conditions (Narrow) - Female




Alcohol-specific mortality - Male




Alcohol-specific mortality - Female




Alcohol-specific mortality - Persons




Mortality from chronic liver disease - Persons




Alcohol-related mortality - Persons



Alcohol-related mortality - Female



Admission episodes for alcohol-related alcoholic liver disease condition (Broad) - Male



Alcohol Related Road Traffic Accidents

